

# Journal Pre-proof

Pediatric Chronic Hand Eczema: Epidemiology, Clinical Presentation and Management Issues

Michael A. Haft, MD, Helen H. Park, MD, Stephanie S. Lee, MD, Jessica M. Sprague, MD, Lawrence F. Eichenfield, MD



PII: S2666-3287(23)00035-4

DOI: <https://doi.org/10.1016/j.jdin.2023.02.008>

Reference: JDIN 341

To appear in: *JAAD International*

Received Date: 14 December 2022

Revised Date: 13 February 2023

Accepted Date: 19 February 2023

Please cite this article as: Haft MA, Park HH, Lee SS, Sprague JM, Eichenfield LF, Pediatric Chronic Hand Eczema: Epidemiology, Clinical Presentation and Management Issues, *JAAD International* (2023), doi: <https://doi.org/10.1016/j.jdin.2023.02.008>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Inc. on behalf of the American Academy of Dermatology, Inc.

1 **Article type:** Review

2 **Title:** Pediatric Chronic Hand Eczema: Epidemiology, Clinical Presentation and Management Issues

3 **Running heading:** Pediatric Chronic Hand Eczema

4 Michael A. Haft, MD<sup>1,2</sup>, Helen H. Park, MD<sup>1,2</sup>, Stephanie S. Lee, MD<sup>1,2</sup>, Jessica M. Sprague, MD<sup>1,2</sup>, Lawrence F.  
5 Eichenfield, MD<sup>1,2</sup>

6 <sup>1</sup>Rady Children's Hospital, San Diego, Division of Pediatric and Adolescent Dermatology, 3020 Children's Way,  
7 San Diego, CA 92130

8 <sup>2</sup>University of California, San Diego, School of Medicine, Department of Dermatology, 9500 Gilman Drive, MC  
9 0869, La Jolla, CA 92093-0869

10

11 **Manuscript word count (Max is 2500 words):** 2585 words [excluding abstract, abbreviations list, references,  
12 figures, tables]

13 **Abstract word count (Max is 200 words):** 158 words

14 **Capsule Summary Word Count:** 49 words

15 **References:** 62

16 **Figures:** 0

17 Supplementary figures: 0

18 **Tables:** 4

19 Supplementary tables: 0

20 Attachments: Cover Letter

21 **Funding:** Internal funding from the University of California, San Diego's Department of Pediatric Dermatology is  
22 funding the publication of this investigation.

23

24 **Statements and Declarations:**

25 **Declaration of interest:**

26 Lawrence F. Eichenfield, MD has served as an advisory board member, and/or speaker, consultant, or clinical trial  
27 investigator for AbbVie, Arcutis, Almirall, Arena, Aslan, Bausch, Castle, Dermavant, Eli Lilly, Forte, Galderma,  
28 Gelnmark/Ichnos, Incyte, Leo Pharma, Novartis, Pfizer, Regeneron, Sanofi Genzyme, UCB and Valeant/Ortho

29 Derm. Michael A. Haft, MD, Helen H. Park, MD, Stephanie S. Lee, MD, and Jessica M. Sprague, MD declare that  
30 they have no conflict of interest.

31 **Availability of data and material:** Data sharing not applicable to this article as no datasets were generated or  
32 analyzed during the current study.

33 **Code availability:** Not applicable

34 **Authors' contributions:** Conceptualization: Michael A. Haft, Lawrence F. Eichenfield; Methodology: Michael A.  
35 Haft, Helen H. Park, Stephanie S. Lee, Jessica M. Sprague, Lawrence F. Eichenfield; Formal analysis and  
36 investigation: Michael A. Haft, Helen H. Park, Stephanie S. Lee, Jessica M. Sprague, Lawrence F. Eichenfield;  
37 Writing - original draft preparation: Michael A. Haft; Writing - review and editing: Michael A. Haft, Helen H. Park,  
38 Stephanie S. Lee, Jessica M. Sprague, Lawrence F. Eichenfield; Resources: Lawrence F. Eichenfield; Supervision:  
39 Jessica M. Sprague, Lawrence F. Eichenfield.

40 **Ethics Approval:** Not applicable as the investigation utilized publicly available and published data to generate a  
41 review.

42 **Consent to participate:** Not applicable

43 **Consent for publication:** Not applicable

44 **Reprint requests:** Lawrence F. Eichenfield

45 **Statement on prior presentation:** Neither the article nor its contents have been presented at a prior presentation by  
46 the authors of this article.

47

48 **Corresponding author:**

49 Lawrence F. Eichenfield, MD

50 3020 Children's Way

51 Mail Code 5092

52 San Diego, CA 92130

53 Telephone number: 858-966-1700 ext.224295

54 Email: leichenfield@rchsd.org, tvbeck@rchsd.org

55

56 **Abstract:**

57           Chronic hand eczema is a persistent inflammatory dermatitis that may significantly affect quality of life,  
58 with psychosocial effects, impact on school, work, and leisure activities, influence on socioeconomic status, and  
59 high healthcare costs. Pediatric chronic hand eczema has a high prevalence yet has not been extensively studied in  
60 children and adolescents. There is minimal published data on pediatric chronic hand eczema in North America, and  
61 no specific management guidelines. Limited prevalence data shows broad ranges (0.9-4.4%) in pre-school and  
62 school children, with one study stating up to 10.0% one-year prevalence for ages 16-19 years. Atopic dermatitis and  
63 allergic contact dermatitis appear important in the pathogenesis of this disease process, though there is limited  
64 pediatric data assessing disease associations and no standardized methodology for evaluating this disorder. Given  
65 the potential life-changing consequences of pediatric chronic hand eczema, further research into this disease process  
66 is warranted to help generate best therapeutic practices and minimize this disease process' morbidity into adulthood.

67  
68 **Capsule summary (Max word count of 50 words):**

- 69       • Pediatric chronic hand eczema has a high prevalence yet has not been extensively studied in children and  
70       adolescents with no guidelines on its management
- 71       • This review on pediatric chronic hand eczema highlights major findings in the literature and supports the  
72       need for further investigation into this life-changing disease process

73  
74  
75 **Keywords:** Pediatric, Child, Children, Childhood, Adolescent, Teenage, Teenager, Young, Chronic, Persistent,  
76 Recalcitrant, Hand, Manus, Eczema, Dermatitis, Skin, Review, Literature, Summary, Questions

77

78 **Introduction:**

79           Chronic hand eczema (CHE) is defined by Deipgen et al. as hand eczema with symptoms persisting for  
80 greater than 3 months or with symptoms returning twice or more within 12 months.<sup>1</sup> CHE significantly affects  
81 quality of life, has financial and psychosocial consequences which include job loss and high healthcare burdens, and  
82 can harm self-esteem and interpersonal relationships.<sup>2-4</sup> The published literature on CHE in children and adolescents  
83 is limited. We evaluate and summarize pediatric chronic hand eczema (P-CHE) epidemiology, risk factors, disease  
84 associations, clinical presentations, severity classifications, diagnostic assessment, and therapeutic interventions.  
85 Knowledge gaps that might drive future research are identified.

86  
87 **Materials and methods:**

88           Systematic search of PubMed, Embase and Cochrane databases was performed from inception through  
89 December 5, 2022, for studies utilizing search terms *hand eczema*, *hand dermatitis*, *hand* and *eczema* or *dermatitis*  
90 in children, restricted to English-language articles. Records were screened according to title and abstracts (985  
91 records), duplicates removed (112), and article eligibility determined by including primary literature or review  
92 articles, observational or controlled studies, scope including patients aged 0-20 years, and a diagnosis of hand  
93 dermatitis or eczema, yielding 31 manuscripts.

94  
95 **Prevalence in Children and Adolescents:**

96           Multiple investigations found that P-CHE is common, with lifetime prevalence of 6.5-13.3% and a 1-year  
97 prevalence of 5.2-10.0% (Table I).<sup>5-10</sup> Figures of prevalence in children vary, with lower prevalence at younger ages.  
98 Wang et al. found that the median age of first occurrence of hand eczema in children was 12 years.<sup>10</sup> Grönhagen et  
99 al. reported hand eczema incidence rates of approximately 0.9% per year in children ages 0-5 and 6-11 years old and  
100 approximately 1.6% in children ages 12-16 years old.<sup>6</sup> Yngveson et al. reported the point prevalence of hand eczema  
101 to be 3.9% (95% CI: 2.9-5.0%) in grade 1 students and 4.4% (95% CI: 3.3-5.5%) in grade 3 students in Sweden,  
102 though differences in data sets were not statistically significant.<sup>9</sup> Much lower childhood lifetime prevalence data was  
103 reported by Crane et al (0.012%), but the Depigen CHE diagnostic criteria were not utilized.<sup>11</sup> Overall prevalence  
104 data correlates with general population data in both children and adults that demonstrates a lifetime prevalence of  
105 about 15% globally.<sup>12</sup>

106 Most studies report that female children are more affected than males, carrying a life-time prevalence of  
107 11.2-16.2% (vs. 6.3-9.6%) and a 1-year prevalence of 6.4-12.5% (vs 4.0-7.3%) (Table I).<sup>6-10</sup> Two studies have  
108 reported higher rates of adult women developing hand eczema before age 20 years (35% and 50%) than men (27%  
109 and 42%),<sup>13,14</sup> and Röhrl and Stenberg observed a positive relationship between hand eczema and female sex (Table  
110 II).<sup>15</sup> However, two recent robust investigations did not find statistically significant odds of P-CHE by sex (Table  
111 II).<sup>10,16</sup> Conflicting sex-difference data in children contrasts with sex-stratified broader population data  
112 demonstrating greater prevalence in the female population.<sup>12</sup>

113 Although these analyses utilize large subject populations, these investigations carry a number of issues,  
114 including recall bias, overestimating the number of participants with hand eczema, and limiting most data sets to one  
115 city in one European nation.<sup>5-10,13-15</sup> Thus, they may not reflect P-CHE's global epidemiology.

116

#### 117 **Symptoms, Distribution, and Morphology:**

118 Symptoms and signs of CHE include itch, redness, scaling, oozing, crusting, and burning pain.<sup>17-19</sup> While  
119 symptoms may be important in assessing P-CHE severity and as potentially measurable parameters in clinical  
120 studies, there is minimal data on signs and symptoms in children and adolescents. Mortz et al. found that children  
121 with hand eczema commonly report pruritus (82.7%), erythema (62.4%), and dry skin with scaling (54.1%).<sup>8</sup>  
122 Simonsen et al. found 26.2% of parents of children with hand eczema reported moderate to severe burning of the  
123 hands, 23.2% with moderate to severe pruritis, 12.6% with moderate to severe pain, and 6.5% with sleep  
124 disturbance.<sup>20</sup> Five investigations reviewed the distribution of lesions in children with hand eczema and found areas  
125 of involvement to be variable. Depending on the study, the most commonly reported locations were dorsal hands,<sup>5</sup>  
126 finger webs or fingers,<sup>8,21</sup> palms,<sup>22</sup> fingertips, or diffuse hand involvement (Table III).<sup>23</sup> Small data sets and reliance  
127 on self-reporting may explain inconsistencies between studies.

128 Few investigations assessed P-CHE severity. One study of 133 children who reported symptoms of hand  
129 eczema in the last 12 months found that 44% lacked any signs or symptoms at the time of evaluation, whereas 13%  
130 had moderate disease and 14% severe disease utilizing the Hand Eczema Extent Score (HEES). Researchers did not  
131 specify how many investigators examined the hands of participants, resulting in possible measurement bias.<sup>6</sup>  
132 Another study of 9 children with P-CHE found an average Hand Eczema Severity Index (HECSI) score correlating

133 to severe disease prior to initiation of alitretinoin therapy, but there was selection bias as all children failed multiple  
134 therapies before starting alitretinoin.<sup>24,25</sup>

135         These 6 studies constitute the bulk of the literature on the signs, symptoms, disease course and outcomes of  
136 in pediatrics. Hand eczema studies in adults show that the disease process presents with edema, erythema, or  
137 vesiculation in its acute form, and fissuring, scaling, or crusting chronically.<sup>19</sup> In a twin cohort study based in  
138 Denmark, of those with hand eczema, 52.3% reported scaling, 51.4% reported erythema, 29.7% reported fissuring,  
139 and 20.7% reported vesicles. Of the 77 adults clinically examined, 47.7% had findings on the fingers (excluding  
140 fingertips), 35.1% on the palms of hands, 30.6% on fingertips, and 24.3% on dorsal hands.<sup>26</sup> Comparisons of these  
141 adult and pediatric data sets are insufficient given the small data sets and reliance on self-reporting in some  
142 investigations.

143         Unanswered questions remain regarding P-CHE's presentation. Like the data available on prevalence, the  
144 published data is limited to Northern Europe. Course and symptom complex are not well categorized in the pediatric  
145 population. Further research into signs and symptoms of this disease presentation in children and adolescents would  
146 be useful.

147

#### 148 **Risk Factors and Diagnoses:**

149         Multiple studies suggest that CHE is strongly associated with atopic dermatitis (AD) in children. In one  
150 study, 43.7% of 0-2 year olds and 54.1% of 3-12 year olds with AD had hand eczema, though chronicity or duration  
151 were not noted.<sup>27</sup> Two investigations from Mortz et al. and one from Grönhagen et al. found odds ratios of 3.7-5.61  
152 between childhood hand eczema and AD (Table II).<sup>8,28,29</sup> In an evaluation of pediatric patients referred for patch  
153 testing to the North American Contact Dermatitis Group, children with hand eczema were more likely to have a  
154 diagnosis of AD than adults.<sup>16</sup> Mortz et al. in 2015 found an odds ratio of 4.3 between hand eczema in childhood  
155 and persistent AD in adulthood,<sup>30</sup> while Wang et al. in 2021 calculated an adjusted odds ratio of 1.8 between  
156 previous diagnosis of AD and lifetime incidence of hand eczema in 15 year olds.<sup>10</sup> Data regarding AD age of onset  
157 and hand eczema risk is conflicting. Wang et al. reported a statistically significant adjusted odds ratio of 1.8 between  
158 early age of onset of AD, independent of the diagnosis of AD itself, and pediatric hand eczema.<sup>10</sup> However,  
159 Grönhagen et al. found no differences between odds ratios of hand eczema and AD at different onset ages of AD.<sup>28</sup>  
160 With regards to hand eczema's relationship to generalized eczema, Silverberg et al. found that hand eczema was

161 associated with lower proportions of generalized dermatitis.<sup>16</sup> Although filaggrin mutations are believed to be  
162 among the strongest risk factors for developing AD,<sup>31</sup> a logistic regression analysis performed by Lagrelius et al.  
163 found no statistically significant odds ratio between filaggrin gene mutations and P-CHE.<sup>32</sup>

164 The data on the relationship of inhalant allergy to pediatric hand eczema is inconsistent. Röhrl and Stenberg  
165 found significant associations between hand eczema and asthma as well as hand eczema and allergic  
166 rhinoconjunctivitis (Table II), but memory bias and use of invalidated questions in this investigation may have  
167 skewed results.<sup>15</sup> Four other investigations found no such links (Table II).<sup>8,10,16,28</sup> Given this discrepant data, these  
168 relationships must be further investigated.

169 P-CHE may also be associated with allergic contact dermatitis (ACD).<sup>29,33</sup> One investigation of children  
170 with AD found that 43.8% of children with hand and/or foot eczema had contact allergy versus 16.0% of children  
171 without hand or foot dermatitis.<sup>33</sup> Another study found that 35.7% of patients with ACD had hand involvement.<sup>34</sup>  
172 Patch testing of children with hand eczema reveals that the most common or most relevant allergens associated with  
173 the disorder include nickel, methylchloroisothiazolinone (MCI) / methylisothiazolinone (MI) (which are commonly  
174 found in cosmetic, hygiene, and household products), and cobalt.<sup>16,22,35-40</sup> Nickel and MI sensitization stand out as  
175 major risk factors for P-CHE, with other allergens less common. Adult population data carries similar findings, as  
176 one report found the most frequent sensitizers in adults with hand eczema to be nickel, MCI/MI, cobalt chloride, and  
177 fragrance mix I.<sup>41</sup>

178 The evidence of irritant contact dermatitis' (ICD) influence on P-CHE is less clear. In 2020 and 2021,  
179 Simonsen et al. found that 26.2% of 0-7 year olds and 36.3% of 5-13 year olds investigated developed hand eczema  
180 following strict hand hygiene protocols upon return to daycare or school in the middle of the COVID-19 pandemic,  
181 with frequency of handwashing, female gender, and history of atopic dermatitis associated with increased risk of  
182 developing hand eczema.<sup>20,42</sup> However, in a 2017 study by Meding et al., investigators found no association between  
183 pediatric hand eczema and hand-water exposure.<sup>43</sup>

184 P-CHE has several coupled diagnoses. Two studies found the most common final diagnoses of children  
185 with P-CHE to be ACD, AD, and vesicular (dyshidrotic) eczema (Table IV).<sup>22,23</sup> Another found the most common  
186 diagnoses for children with CHE referred for patch testing to be ACD, AD, and ICD (Table IV).<sup>16</sup> This suggests that  
187 ACD and AD are commonly associated with CHE in childhood. The adult CHE literature presents some overlap in  
188 findings, with one analysis presenting the most common associated diagnoses as combinations of ICD, ACD, and



189 vesicular eczema,<sup>1</sup> suggesting that AD plays a greater role in P-CHE pathogenesis with ICD playing a greater role in  
190 adult CHE. In clinical practice, it appears that some children and adolescents have significant chronic hand  
191 dermatitis as part of a constellation of findings in active AD, while others have localized CHE, or predominate  
192 issues with CHE disproportionate to other issues with AD. We believe the term CHE remains useful, with  
193 subcategories of etiology including AD and ACD.

194         Investigations show conflicting data regarding the influence of inhalant allergy and ICD on P-CHE. AD  
195 and ACD's overlap with and impact on P-CHE are much clearer,<sup>16</sup> and evidence demonstrates nickel and MI  
196 allergy's influence on hand eczema in children. Further investigations need to elucidate the relationship of these and  
197 other risk factors for the development and persistence of CHE in childhood.

198         Many methods of diagnosis/classification of CHE<sup>44-47</sup> attempt to incorporate various combinations of  
199 morphology, etiology, and chronological progression, while major studies have found insignificant association  
200 between classification and etiology.<sup>45,46,48</sup>

201

202

### 203 **Diagnostic Testing, Severity Assessment, and Therapeutics:**

204         P-CHE workup frequently includes patch testing with studies finding that anywhere from 14.5-28.0% of  
205 children referred for patch testing have hand eczema.<sup>16,22,35,49</sup> In two studies, patch testing was reported to have a  
206 clinical relevance of 78% in P-CHE and 76.2% in pediatric hand eczema,<sup>22,50</sup> much higher than in adult studies.<sup>51,52</sup>  
207 The literature has not supported IgE testing as no association has been found between positive specific IgE during  
208 childhood and P-CHE.<sup>28</sup>

209         The use of standardized severity assessments are rare in the P-CHE literature, with one P-CHE study  
210 utilizing the HECSI and Investigator Global Assessment (IGA)<sup>24</sup> and one study utilizing the HEES.<sup>6</sup> Other  
211 evaluation measures, including the Dermatology Life Quality Index (DLQI),<sup>17</sup> Quality of Life in Hand Eczema  
212 Questionnaire (QOLHEQ),<sup>53</sup> and modified total lesion symptom score (mTLSS)<sup>54</sup> appear to only be executed in  
213 adult populations or in studies containing mixed populations of both children and adults.<sup>55</sup>

214         Studies evaluating topical or systemic medications for P-CHE are limited. In a retrospective analysis of 13  
215 children who received systemic alitretinoin therapy, 9 were children with CHE. In this subgroup, 7/9 had moderate  
216 to excellent results on alitretinoin.<sup>24</sup> In retrospective review of 75 children receiving phototherapy for cutaneous

217 conditions, 4 had severe hand eczema, of which 3 had clinical improvement after psoralen and ultraviolet A (PUVA)  
218 therapy.<sup>56</sup> In a 2019 systematic review of publications on hand eczema therapeutics performed by Christoffers et al.,  
219 researchers could not find a single study on therapeutics exclusively in pediatrics. Most of the studies excluded  
220 children and pediatric patients were not given their own subgroup analysis apart from adults in any article.<sup>57</sup>

221 Although the literature lacks published data on P-CHE treatment, investigators from this article and the  
222 Pediatric Dermatology Research Alliance, performed a survey of pediatric dermatologist CHE experts. Surveyed  
223 respondents all utilize topical corticosteroids (TCS) as first line topical therapy with most choosing TCS, topical  
224 calcineurin inhibitors, and topical phosphodiesterase-4 inhibitors as second line agents. Systemic treatment use is  
225 rare, with most respondents reporting 5 or fewer patients treated for the indication of P-CHE. The most preferred  
226 systemic agent for P-CHE was dupilumab, followed by methotrexate.<sup>58</sup>

227 No specific guidelines exist for P-CHE management, though there are published guidelines and consensus  
228 statements for the management of CHE based on adult data.<sup>59,60</sup> The European Society of Contact Dermatitis  
229 (ESCD) produced updated management guidelines for hand eczema in 2022, recommending the use of patch testing  
230 in all patients with CHE. Other recommendations included skin prick testing, microbial testing, and cutaneous  
231 biopsy when deemed appropriate. However, there is significant disagreement among experts concerning utility of  
232 patch testing irrespective of morphology and location, predictive value of testing, and cost effectiveness.<sup>48</sup>  
233 Management includes prevention and use of therapeutics from emollients and topical steroids to systemic agents  
234 such as oral alitretinoin (approved for CHE in Europe and the United Kingdom) or cyclosporin.<sup>61</sup> Recent literature  
235 highlights the use of emerging and investigational systemic agents including biologic agents and JAK inhibitors for  
236 CHE in adults.<sup>62</sup>

237 The lack of scoring systems, published data on therapeutics, and management guidelines focused on the  
238 pediatric population is troubling given the life-altering potential of this disorder.<sup>4</sup> Utilization of standardized metrics  
239 of disease severity, quality of life, and treatment response could assist in determining the comparative efficacy of  
240 various interventions and guide the development of best practices guidelines.

241

#### 242 **Future Directions and Significance of Further Investigations:**

243 There remain wide knowledge gaps in the epidemiology, presentation, risk stratification, diagnosis, and  
244 management of CHE in pediatric populations. Most published studies are limited to patients in Northern Europe.

245 Most of the data on CHE epidemiology is based on adult patients.<sup>12</sup> While many adults report hand eczema onset in  
246 childhood, few reports investigate characteristics of this disease in the pediatric population, including assessing what  
247 percentage of children with hand eczema have AD. Even fewer studies have explored scoring systems in assessment  
248 or therapeutic management of P-CHE.

249 Numerous questions remain in all domains of this disorder in children and adolescents: What is the course  
250 of hand eczema in childhood versus adults? How does hand eczema in pediatrics progress from acute to chronic  
251 disease? What percentage of those affected have active AD or other inflammatory skin conditions? What are other  
252 risk factors for disease development? What classification systems are ideal? Why is there little consensus on features  
253 and testing? When would providers utilize certain tests for disease workup and treatments for disease management?  
254 How do clinicians assess treatment response? How does therapeutic data correlate with severity or quality of life  
255 scoring?

256 Early recognition and treatment of this disease process in childhood may minimize the disease impact,  
257 decrease healthcare burden, and improve quality of life.<sup>4</sup> Further investigations into the epidemiology of P-CHE  
258 onset and course, disease associations, comorbidities, and therapeutics are important to determine best practices to  
259 allow for comprehensive and successful management. With ongoing development of new topical and systemic  
260 agents for CHE as well as for AD, focused research on P-CHE is warranted.

261

262

263 **Abbreviations and Acronyms:**

264 ACD: Allergic Contact Dermatitis

265 AD: Atopic Dermatitis

266 CHE: Chronic Hand Eczema

267 CI: Confidence Interval

268 DLQI: Dermatology Life Quality Index

269 ESCD: European Society of Contact Dermatitis

270 HE: Hand Eczema

271 HECSI: Hand Eczema Severity Index

272 HEES: Hand Eczema Extent Score

273 ICD: Irritant Contact Dermatitis

- 274 IGA: Investigator Global Assessment
- 275 MCI: Methylchloroisothiazolinone
- 276 MI: Methylisothiazolinone
- 277 NR: Not Reported
- 278 mTLSS: Modified Total Lesion Symptom Score
- 279 P-CHE: Pediatric Chronic Hand Eczema
- 280 PPD: Paraphenylenediamine
- 281 PUVA: Psoralen and Ultraviolet A
- 282 QOLHEQ: Quality of Life in Hand Eczema Questionnaire
- 283
- 284 **Acknowledgements:** None
- 285

286 **References**

- 287 1. Diepgen TL, Elsner P, Schliemann S, et al. Guideline on the management of hand eczema  
288 ICD-10 Code: L20. L23. L24. L25. L30. *J Dtsch Dermatol Ges.* May 2009;7 Suppl 3:S1-16.  
289 doi:10.1111/j.1610-0387.2009.07061.x
- 290 2. Armstrong A, Hahn-Pedersen J, Bartlett C, Glanville J, Thyssen JP. Economic Burden of  
291 Chronic Hand Eczema: A Review. *Am J Clin Dermatol.* Mar 8 2022;doi:10.1007/s40257-021-  
292 00669-6
- 293 3. Ahmed A, Shah R, Papadopoulos L, Bewley A. An ethnographic study into the  
294 psychological impact and adaptive mechanisms of living with hand eczema. *Clin Exp Dermatol.*  
295 Jul 2015;40(5):495-501. doi:10.1111/ced.12619
- 296 4. Cazzaniga S, Ballmer-Weber BK, Grani N, et al. Medical, psychological and socio-  
297 economic implications of chronic hand eczema: a cross-sectional study. *J Eur Acad Dermatol*  
298 *Venereol.* Apr 2016;30(4):628-37. doi:10.1111/jdv.13479
- 299 5. Dotterud LK, Falk ES. Contact allergy in relation to hand eczema and atopic diseases in  
300 north Norwegian schoolchildren. *Acta Paediatr.* Apr 1995;84(4):402-6. doi:10.1111/j.1651-  
301 2227.1995.tb13659.x
- 302 6. Gronhagen CM, Liden C, Bergstrom A, Kull I, Wahlgren CF, Meding B. Prevalence and  
303 incidence of hand eczema in adolescence: report from BAMSE--a population-based birth cohort.  
304 *Br J Dermatol.* Sep 2014;171(3):609-14. doi:10.1111/bjd.13194
- 305 7. Johannisson A, Ponten A, Svensson A. Prevalence, incidence and predictive factors for  
306 hand eczema in young adults - a follow-up study. *BMC Dermatol.* Oct 29 2013;13:14.  
307 doi:10.1186/1471-5945-13-14
- 308 8. Mortz CG, Lauritsen JM, Bindslev-Jensen C, Andersen KE. Prevalence of atopic  
309 dermatitis, asthma, allergic rhinitis, and hand and contact dermatitis in adolescents. The Odense  
310 Adolescence Cohort Study on Atopic Diseases and Dermatitis. *Br J Dermatol.* Mar  
311 2001;144(3):523-32. doi:10.1046/j.1365-2133.2001.04078.x
- 312 9. Yngveson M, Svensson A, Isacson A. Prevalence of self-reported hand dermatosis in  
313 upper secondary school pupils. *Acta Derm Venereol.* Sep 1998;78(5):371-4.  
314 doi:10.1080/000155598443097
- 315 10. Wang J, Tischer C, Standl M, et al. Lifetime prevalence and determinants of hand eczema  
316 in an adolescent population in Germany: 15-year follow-up of the LISA cohort study. *J Eur Acad*  
317 *Dermatol Venereol.* Nov 15 2021;doi:10.1111/jdv.17814
- 318 11. Crane MM, Webb DJ, Watson E, Cunliffe T, English J. Hand eczema and steroid-  
319 refractory chronic hand eczema in general practice: prevalence and initial treatment. *Br J*  
320 *Dermatol.* Apr 2017;176(4):955-964. doi:10.1111/bjd.14974
- 321 12. Thyssen JP, Johansen JD, Linneberg A, Menné T. The epidemiology of hand eczema in  
322 the general population--prevalence and main findings. *Contact Dermatitis.* Feb 2010;62(2):75-  
323 87. doi:10.1111/j.1600-0536.2009.01669.x
- 324 13. Meding B, Järvalho B. Incidence of hand eczema-a population-based retrospective  
325 study. *J Invest Dermatol.* Apr 2004;122(4):873-7. doi:10.1111/j.0022-202X.2004.22406.x
- 326 14. Vindenes HK, Svanes C, Lygre SHL, Hollund BE, Langhammer A, Bertelsen RJ.  
327 Prevalence of, and work-related risk factors for, hand eczema in a Norwegian general population  
328 (The HUNT Study). Article. *Contact Dermatitis.* 2017;77(4):214-223. doi:10.1111/cod.12800
- 329 15. Röhrh K, Stenberg B. Lifestyle factors and hand eczema in a Swedish adolescent  
330 population. *Contact Dermatitis.* Mar 2010;62(3):170-6. doi:10.1111/j.1600-0536.2009.01679.x

- 331 16. Silverberg JI, Warshaw EM, Maibach HI, et al. Hand Eczema in Children Referred for  
332 Patch Testing: North American Contact Dermatitis Group Data, 2000-2016. *Br J Dermatol*. Jan  
333 16 2021;doi:10.1111/bjd.19818
- 334 17. Barrett A, Hahn-Pedersen J, Kragh N, Evans E, Gnanasakthy A. Patient-Reported  
335 Outcome Measures in Atopic Dermatitis and Chronic Hand Eczema in Adults. *Patient*. Oct  
336 2019;12(5):445-459. doi:10.1007/s40271-019-00373-y
- 337 18. Rademaker M, Armour K, Baker C, et al. Management of chronic hand and foot eczema.  
338 An Australia/New Zealand Clinical narrative. *Australas J Dermatol*. Feb 2021;62(1):17-26.  
339 doi:10.1111/ajd.13418
- 340 19. Agner T, Elsner P. Hand eczema: epidemiology, prognosis and prevention. *J Eur Acad*  
341 *Dermatol Venereol*. Jan 2020;34 Suppl 1:4-12. doi:10.1111/jdv.16061
- 342 20. Simonsen AB, Ruge IF, Quaade AS, Johansen JD, Thyssen JP, Zachariae C. Increased  
343 occurrence of hand eczema in young children following the Danish hand hygiene  
344 recommendations during the COVID-19 pandemic. Article. *Contact Dermatitis*. 2021;84(3):144-  
345 152. doi:10.1111/cod.13727
- 346 21. Lee HJ, Ha SJ, Ahn WK, et al. Clinical evaluation of atopic hand-foot dermatitis. *Pediatr*  
347 *Dermatol*. Mar-Apr 2001;18(2):102-6. doi:10.1046/j.1525-1470.2001.018002102.x
- 348 22. Toledo F, García-Bravo B, Fernández-Redondo V, et al. Patch testing in children with  
349 hand eczema. A 5-year multicentre study in Spain. *Contact Dermatitis*. Oct 2011;65(4):213-9.  
350 doi:10.1111/j.1600-0536.2011.01943.x
- 351 23. Ortiz-Salvador JM, Subiabre-Ferrer D, Garcia Rabasco A, Esteve-Martinez A, Zaragoza-  
352 Ninet V, Alegre de Miquel V. [Hand eczema in children. Clinical and epidemiological study of  
353 the population referred to a tertiary hospital]. *An Pediatr (Barc)*. Jun 2018;88(6):309-314.  
354 Eccema de manos en niños. Estudio clínico-epidemiológico de la población remitida a un  
355 hospital terciario. doi:10.1016/j.anpedi.2017.07.005
- 356 24. Luchsinger I, Vogler T, Schwieger-Briel A, et al. Safe and effective use of alitretinoin in  
357 children with recalcitrant hand eczema and other dermatoses - a retrospective analysis. *J Eur*  
358 *Acad Dermatol Venereol*. May 2020;34(5):1037-1042. doi:10.1111/jdv.16088
- 359 25. Oosterhaven JAF, Schuttelaar MLA. Responsiveness and interpretability of the Hand  
360 Eczema Severity Index. *Br J Dermatol*. Apr 2020;182(4):932-939. doi:10.1111/bjd.18295
- 361 26. Lerbaek A, Kyvik KO, Ravn H, Menne T, Agner T. Clinical characteristics and  
362 consequences of hand eczema - an 8-year follow-up study of a population-based twin cohort.  
363 *Contact Dermatitis*. Apr 2008;58(4):210-6. doi:10.1111/j.1600-0536.2007.01305.x
- 364 27. Simpson EL, Thompson MM, Hanifin JM. Prevalence and morphology of hand eczema  
365 in patients with atopic dermatitis. *Dermatitis*. Sep 2006;17(3):123-7.  
366 doi:10.2310/6620.2006.06005
- 367 28. Grönhagen C, Lidén C, Wahlgren CF, et al. Hand eczema and atopic dermatitis in  
368 adolescents: a prospective cohort study from the BAMSE project. *Br J Dermatol*. Nov  
369 2015;173(5):1175-82. doi:10.1111/bjd.14019
- 370 29. Mortz CG, Lauritsen JM, Bindslev-Jensen C, Andersen KE. Contact allergy and allergic  
371 contact dermatitis in adolescents: prevalence measures and associations. The Odense  
372 Adolescence Cohort Study on Atopic Diseases and Dermatitis (TOACS). *Acta Derm Venereol*.  
373 2002;82(5):352-8. doi:10.1080/000155502320624087
- 374 30. Mortz CG, Andersen KE, Dellgren C, Barington T, Bindslev-Jensen C. Atopic dermatitis  
375 from adolescence to adulthood in the TOACS cohort: prevalence, persistence and comorbidities.  
376 *Allergy*. Jul 2015;70(7):836-45. doi:10.1111/all.12619

- 377 31. O'Regan GM, Sandilands A, McLean WHI, Irvine AD. Filaggrin in atopic dermatitis. *J*  
378 *Allergy Clin Immunol*. Oct 2008;122(4):689-693. doi:10.1016/j.jaci.2008.08.002
- 379 32. Lagrelus M, Wahlgren CF, Bradley M, et al. Filaggrin gene mutations in relation to  
380 contact allergy and hand eczema in adolescence. *Contact Dermatitis*. Mar 2020;82(3):147-152.  
381 doi:10.1111/cod.13444
- 382 33. Isaksson M, Olhardt S, Radehed J, Svensson A. Children with Atopic Dermatitis Should  
383 Always be Patch-tested if They Have Hand or Foot Dermatitis. *Acta Derm Venereol*. May  
384 2015;95(5):583-6. doi:10.2340/00015555-1995
- 385 34. Yilmaz Z, Ozkaya E. Patch-test results in terms of the recently recommended allergens in  
386 children and adolescents: A retrospective cohort study over 22 years from Turkey. *Contact*  
387 *Dermatitis*. Mar 23 2021;doi:10.1111/cod.13842
- 388 35. Beattie PE, Green C, Lowe G, Lewis-Jones MS. Which children should we patch test?  
389 *Clin Exp Dermatol*. Jan 2007;32(1):6-11. doi:10.1111/j.1365-2230.2006.02232.x
- 390 36. Mortz CG, Lauritsen JM, Bindslev-Jensen C, Andersen KE. Nickel sensitization in  
391 adolescents and association with ear piercing, use of dental braces and hand eczema. The Odense  
392 Adolescence Cohort Study on Atopic Diseases and Dermatitis (TOACS). *Acta Derm Venereol*.  
393 2002;82(5):359-64. doi:10.1080/000155502320624096
- 394 37. Urwin R, Wilkinson M. Methylchloroisothiazolinone and methylisothiazolinone contact  
395 allergy: a new 'epidemic'. *Contact Dermatitis*. Apr 2013;68(4):253-5. doi:10.1111/cod.12064
- 396 38. Geier J, Lessmann H, Schnuch A, Uter W. Recent increase in allergic reactions to  
397 methylchloroisothiazolinone/methylisothiazolinone: is methylisothiazolinone the culprit?  
398 *Contact Dermatitis*. Dec 2012;67(6):334-41. doi:10.1111/j.1600-0536.2012.02157.x
- 399 39. Lundov MD, Thyssen JP, Zachariae C, Johansen JD. Prevalence and cause of  
400 methylisothiazolinone contact allergy. *Contact Dermatitis*. Sep 2010;63(3):164-7.  
401 doi:10.1111/j.1600-0536.2010.01774.x
- 402 40. Mowad CM. Methylchloro-isothiazolinone revisited. *Am J Contact Dermat*. Jun  
403 2000;11(2):115-8. doi:10.1053/ac.2000.6344
- 404 41. Boonstra MB, Christoffers WA, Coenraads PJ, Schuttelaar ML. Patch test results of hand  
405 eczema patients: relation to clinical types. *J Eur Acad Dermatol Venereol*. May 2015;29(5):940-  
406 7. doi:10.1111/jdv.12735
- 407 42. Simonsen AB, Ruge IF, Quaade AS, Johansen JD, Thyssen JP, Zachariae C. High  
408 incidence of hand eczema in Danish schoolchildren following intensive hand hygiene during the  
409 COVID-19 pandemic: a nationwide questionnaire study. *Br J Dermatol*. Nov 2020;183(5):975-  
410 976. doi:10.1111/bjd.19413
- 411 43. Meding B, Gronhagen CM, Bergstrom A, Kull I, Wrangsjo K, Liden C. Water Exposure  
412 on the Hands in Adolescents: A Report from the BAMSE Cohort. *Acta Derm Venereol*. Feb 8  
413 2017;97(2):188-192. doi:10.2340/00015555-2505
- 414 44. Agner T, Aalto-Korte K, Andersen KE, et al. Classification of hand eczema. *J Eur Acad*  
415 *Dermatol Venereol*. Dec 2015;29(12):2417-22. doi:10.1111/jdv.13308
- 416 45. Diepgen TL, Andersen KE, Brandao FM, et al. Hand eczema classification: a cross-  
417 sectional, multicentre study of the aetiology and morphology of hand eczema. *Br J Dermatol*.  
418 Feb 2009;160(2):353-8. doi:10.1111/j.1365-2133.2008.08907.x
- 419 46. Johansen JD, Hald M, Andersen BL, et al. Classification of hand eczema: clinical and  
420 aetiological types. Based on the guideline of the Danish Contact Dermatitis Group. *Contact*  
421 *Dermatitis*. Jul 2011;65(1):13-21. doi:10.1111/j.1600-0536.2011.01911.x

- 422 47. Molin S, Diepgen TL, Ruzicka T, Prinz JC. Diagnosing chronic hand eczema by an  
423 algorithm: a tool for classification in clinical practice. *Clin Exp Dermatol*. Aug 2011;36(6):595-  
424 601. doi:10.1111/j.1365-2230.2010.04001.x
- 425 48. Silverberg JI, Guttman-Yassky E, Agner T, et al. Chronic Hand Eczema Guidelines From  
426 an Expert Panel of the International Eczema Council. *Dermatitis*. Sep 30  
427 2020;doi:10.1097/DER.0000000000000659
- 428 49. Simonsen AB, Deleuran M, Mortz CG, Johansen JD, Sommerlund M. Allergic contact  
429 dermatitis in Danish children referred for patch testing - a nationwide multicentre study. *Contact*  
430 *Dermatitis*. Feb 2014;70(2):104-11. doi:10.1111/cod.12129
- 431 50. Toledo Alberola F, Silvestre Salvador JF, Latorre Martínez N, Cuesta Montero L,  
432 Monteagudo Paz A, Betlloch Mas I. Utility of patch-tests in childrens hand eczema: Comparative  
433 study with eczema in the pediatric population and with adult hand eczema. Conference Abstract.  
434 *Contact Dermatitis*. 2010;63:68. doi:10.1111/j.1600-0536.2010.01795.x
- 435 51. Minh PPT, Minh TT, Huu DL, et al. Using Patch Testing to Improve Therapeutic  
436 Outcome in the Treatment of Hand Eczema in Vietnamese Patients. *Open Access Maced J Med*  
437 *Sci*. Jan 30 2019;7(2):204-207. doi:10.3889/oamjms.2019.051
- 438 52. Agner T, Andersen KE, Brandao FM, et al. Contact sensitisation in hand eczema  
439 patients-relation to subdiagnosis, severity and quality of life: a multi-centre study. *Contact*  
440 *Dermatitis*. Nov 2009;61(5):291-6. doi:10.1111/j.1600-0536.2009.01630.x
- 441 53. Ofenloch RF, Weisshaar E, Dumke AK, Molin S, Diepgen TL, Apfelbacher C. The  
442 Quality of Life in Hand Eczema Questionnaire (QOLHEQ): validation of the German version of  
443 a new disease-specific measure of quality of life for patients with hand eczema. *Br J Dermatol*.  
444 Aug 2014;171(2):304-12. doi:10.1111/bjd.12819
- 445 54. Shroff A, Malajian D, Czarnowicki T, et al. Use of 308 nm excimer laser for the  
446 treatment of chronic hand and foot eczema. *Int J Dermatol*. Aug 2016;55(8):e447-53.  
447 doi:10.1111/ijd.13205
- 448 55. Ronsch H, Apfelbacher C, Brans R, et al. Which outcomes have been measured in hand  
449 eczema trials? A systematic review. *Contact Dermatitis*. Apr 2019;80(4):201-207.  
450 doi:10.1111/cod.13212
- 451 56. Eustace K, Dolman S, Alsharqi A, Sharpe G, Parslew R. Use of Phototherapy in  
452 Children. *Pediatr Dermatol*. Mar 2017;34(2):150-155. doi:10.1111/pde.13072
- 453 57. Christoffers WA, Coenraads PJ, Svensson A, et al. Interventions for hand eczema.  
454 *Cochrane Database Syst Rev*. Apr 26 2019;4:CD004055.  
455 doi:10.1002/14651858.CD004055.pub2
- 456 58. Data presented at the International Society of Atopic Dermatitis 2022 Meeting, October  
457 18, 2022.
- 458 59. English J, Aldridge R, Gawkrödger DJ, et al. Consensus statement on the management of  
459 chronic hand eczema. *Clin Exp Dermatol*. Oct 2009;34(7):761-9. doi:10.1111/j.1365-  
460 2230.2009.03649.x
- 461 60. Silvestre Salvador JF, Heras Mendaza F, Hervella Garces M, et al. Guidelines for the  
462 Diagnosis, Treatment, and Prevention of Hand Eczema. *Actas Dermosifiliogr*. Jan 2  
463 2020;111(1):26-40. Guia para el diagnostico, el tratamiento y la prevencion del eccema de  
464 manos. doi:10.1016/j.ad.2019.04.005
- 465 61. Thyssen JP, Schuttelaar MLA, Alfonso JH, et al. Guidelines for diagnosis, prevention,  
466 and treatment of hand eczema. *Contact Dermatitis*. May 2022;86(5):357-378.  
467 doi:10.1111/cod.14035



468 62. Cheng J, Facheris P, Ungar B, Guttman-Yassky E. Current emerging and investigational  
469 drugs for the treatment of chronic hand eczema. *Expert Opin Investig Drugs*. Aug  
470 2022;31(8):843-853. doi:10.1080/13543784.2022.2087059  
471

472

Journal Pre-proof

473 **Table legend**474 **Table I.** Study Data on the Prevalence of Hand Eczema in Pediatrics475 **Table II.** Study Data on Risk Factors for Pediatric Hand Eczema476 **Table III.** Study Data on the Physical Distribution of Hand Eczema in Pediatrics477 **Table IV.** Study Data of Diagnoses Associated with Pediatric Hand Eczema

478

479 Table I. "Study Data on the Prevalence of Hand Eczema in Pediatrics"

Source	Setting	Study Design (Years)	Age of Study Participants Included (Years)	No. of Total Participants	No. of Females	No. of Males	No. of Participants Reporting			No. of Females Reporting		No. of Males Reporting	
							Lifetime Prevalence of Hand Eczema (%)	1-year Prevalence of Hand Eczema (%)	Current Hand Eczema (%)	Lifetime Prevalence of Hand Eczema (%)	1-year Prevalence of Hand Eczema (%)	Lifetime Prevalence of Hand Eczema (%)	1-year Prevalence of Hand Eczema (%)
Grönshagen et al., 2014	Sweden; Birth registry	Birth cohort (1994-2012)	0-16	2927	1494	1433	284 (9.7)	152 (5.2)	NR	168 (11.2)	95 (6.4)	116 (8.1)	57 (4.0)
Johannisson et al., 2013	Sweden; 4 schools	Prospective Cohort (1995)	16-19	1516	857	659	202 (13.3)	NR	NR	139 (16.2)	NR	63 (9.6)	NR
Mortz et al., 2001	Denmark; 40 schools	Cross-sectional (1995-1997)	12-16	1438	713	725	133 (9.2)	105 (7.3)	46 (3.2)	87 (12.2)	72 (10.1)	46 (6.3)	33 (4.6)
Wang et al., 2021	Germany; 4 regions	Cross-sectional (2012-2014)	15	1468	715	753	153 (10.4)	NR	NR	91 (12.7)	NR	62 (8.2)	NR
Yngveson et al., 1998	Sweden; 4 schools	Cross-sectional (1995)	16-19	2572	1314	1258	NR	257 (10.0)	108 (4.2)	NR	322 (12.5)	NR	188 (7.3)

NR, Not reported

480

481 Table II. "Study Data on Risk Factors for Pediatric Hand Eczema"

482

Source	Setting	Study Design (Years)	Age of Study Participants Included (Years)	No. of Total Participants	Odds Ratio of Association Between				
					Female Sex and HE (95% CI)	AD and HE (95% CI)	Asthma and HE (95% CI)	Allergic Rhinitis and HE (95% CI)	Nickel Allergy and HE (95% CI)
Grönshagen et al., 2015	Sweden; Birth registry	Birth cohort (1994-2012)	0-16	2927	NR	3.7 (2.0-7.0) (P<0.01)	1.5 (0.8-2.5) (P=0.2), 1.2 (0.6-2.1) (P=0.6) <sup>1</sup>	NR	NR
Mortz et al., 2001	Denmark; 40 schools	Cross-sectional (1995-1997)	12-16	1438	NR	5.61 (3.81-8.25) (P<0.001)	1.58 (1.01-2.46) (P<0.05) (Insignificant after Bonferroni correction) <sup>1</sup>	NR	NR
Röhl and Stenberg, 2010	Sweden; 11 schools	Cross-sectional (2000-2004)	14-24	7543	2.0 (1.3-3.2)	4.5 (3.3-6.1)	1.48 (1.04-2.09)	1.8 (1.3-2.5)	1.7 (1.1-2.7)
Silverberg et al., 2021	USA, Canada; >20 clinics	Retrospective (2000-2016)	0-18	1634	0.525 (0.497-0.554) (P=0.6341)	0.989 (0.884-1.431) (P=0.9550)	0.622 (0.378-1.023) (P=0.0615)	0.782 (0.511-1.197) (P=0.2578)	0.539 (0.349-0.832) (P=0.00525)
Wang et al., 2021	Germany; 4 regions	Cross-sectional (2012-2014)	15	1468	1.5 (0.9-2.6) (P=0.090)	1.8 (1.1-2.8) (P=0.019)	NR <sup>1</sup>	1.4 (0.8-2.5) (P=0.250)	NR

AD, Atopic dermatitis; CI, Confidence interval; HE, Hand eczema; NR, Not reported  
<sup>1</sup>Also included allergic rhinitis in their calculation  
<sup>2</sup>Not reported in multivariable logistic regression analysis

483

484 Table III. "Study Data on the Physical Distribution of Hand Eczema in Pediatrics"

485

Table III: Study Data on the Physical Distribution of Hand Eczema in Pediatrics

Source	Setting	Study Design (Years)	Age of Study Participants Included (Years)	No. of Total Participants	No. of Participants with Lifetime Prevalence of Hand Eczema (%)	Out of All Participants Reporting Lifetime Prevalence of Hand Eczema, No. of Those with				
						Hand Diffusely Affected (%)	Fingers / Finger-webs / Lateral fingers (%)	Dorsal hands (%)	Palms (%)	Fingertips (%)
Dotterud & Falk, 1995	Norway; Multiple schools	Cross-sectional (1995)	7-12	551	36 (6.5)	NR	NR	14 (38.9)	1 (2.8)	NR
Lee et al., 2001	South Korea; 1 hospital	Cross-sectional (1997-1998)	0.5-12	108	62 (57.0)	NR	NR <sup>1</sup>	38 (61.3)	48 (77.4)	NR
Mortz et al., 2001	Denmark; 40 schools	Cross-sectional (1995-1997)	12-16	1438	133 (9.2)	NR	86 (64.7)	68 (51.1)	22 (16.5)	NR
Ortiz-Salvador et al., 2018	Spain; 1 hospital	Retrospective observational (1996-2016)	0-16	389	42 (10.8)	12 (28.6)	4 (9.5)	5 (11.9)	9 (21.4)	12 (28.6)
Toledo et al., 2011	Spain; 11 hospitals	Retrospective multicenter (2005-2009)	0-15	480	111 (23.1)	19 (17.1)	15 (13.5)	7 (6.3)	29 (26.1)	14 (12.6)

NR, Not reported  
<sup>1</sup>Separately reported affected dorsal and ventral surfaces of fingers of right and left hands

486

487

488 Table IV. "Study Data of Diagnoses Associated with Pediatric Hand Eczema"

489

Table IV: Study Data of Diagnoses Associated with Pediatric Hand Eczema

Source	Setting	Study Design (Years)	Age of Study Participants Included (Years)	No. of Total Participants	No. of Participants with Lifetime Prevalence of Hand Eczema (%)	No. of Participants with Hand Eczema Diagnosed with				
						Atopic Dermatitis (%)	Allergic Contact Dermatitis (%)	Irritant Contact Dermatitis (%)	Hyperkeratotic Eczema (%)	Vesicular Eczema (%)
Ortiz-Salvador et al., 2018	Spain; 1 hospital	Retrospective observational (1996-2016)	0-16	389	42 (10.8)	15 (35.7)	14 (33.3)	2 (4.8)	5 (11.9)	6 (14.3)
Silverberg et al., 2021	USA, Canada; >20 clinics	Retrospective (2000-2016)	0-18	1634	237 (14.5)	88 (37.1)	117 (49.4)	40 (16.9)	NR	10 (4.2)
Toledo et al., 2011	Spain; 11 hospitals	Retrospective multicenter (2005-2009)	0-15	480	111 (23.1)	32 (28.8)	40 (36)	17 (15.3)	NR	18 (16.2)

NR, Not reported

490

491

492

1 **Capsule summary (Max word count of 50 words):**

- 2       • Pediatric chronic hand eczema has a high prevalence yet has not been extensively studied in children and  
3       adolescents with no guidelines on its management
- 4       • This review on pediatric chronic hand eczema highlights major findings in the literature and supports the  
5       need for further investigation into this life-changing disease process

6

Journal Pre-proof